

HODGE STRUCTURE OF A COMPLETE INTERSECTION OF QUADRICS IN A PROJECTIVE SPACE

SU-JEONG KANG

Let V be a smooth projective variety of dimension n . One may say that V has motivic dimension less than $d + 1$ if the cohomology of V comes from varieties of dimensions less than $d + 1$ in some geometric way. More precisely, we say that V has *motivic dimension* less than $d + 1$ if there exists a (nonconnected) smooth projective variety W of dimension less than $d + 1$ and an algebraic correspondence Γ on $W \times X$ such that Γ induces a surjection $H^*(W) \rightarrow H^*(V)$.

Suppose the Generalized Hodge conjecture (GHC) ([G],[L]) holds for V and suppose the level of V is less than l (i.e. $\text{level}(H^i(V)) < l$ for all i). This implies that V has a motivic dimension less than l . Conversely, if V has motivic dimension less than $d + 1$, i.e. there is a smooth projective variety W of dimension less than $d + 1$ and a surjection $\phi : H^*(W) \rightarrow H^n(V)$ induced by a correspondence Γ on $W \times X$, then the level of the Hodge structure $H^n(V)$ is less than $d + 1$ because a morphism of Hodge structures preserves the level. The existence of W does not imply that the GHC holds for V of course, because the dimension of the variety W is not small enough to conclude the GHC for V . However, this gives a way to reduce the GHC for V to the GHC for W , the variety with the smaller dimension. And we think that this can be a practical intermediate step for checking GHC for a smooth projective variety.

For a smooth complete intersection of k quadrics $V = Q_1 \cap \cdots \cap Q_k$ in $\mathbb{P}^{n+k}$, it can be checked that the level of the Hodge structure $H^n(V)$ is less than k (for example by checking Hodge number of it). In this paper, we show that a smooth complete intersection of k quadrics in $\mathbb{P}^{n+k}$ has motivic dimension less than k (Theorem 1.2). As a corollary of this, we get the Hodge conjecture for V holds if $k < 4$.

A brief outline of the proof of main theorem is as follows. By using the construction in [O], we form a family of quadrics parametrized by $\mathbb{P}^{k-1}$ with the base locus V . In case when $n+k$ is odd, this is a family of even dimensional quadrics. Then we use the fact that an even dimensional quadric contains two irreducible families of linear spaces of the expected dimension, and we choose W to be a double covering of $\mathbb{P}^{n+k}$. In the case when $n+k$ is even, then quadrics in the original family are odd dimensional. So in order to find W , we pass to the family of singular fibers over the discriminant variety which can be understood as a family of quadrics of even dimension. Then we form a double covering of the discriminant variety to find W .

All varieties in this paper will be defined over $\mathbb{C}$ and the cohomology without coefficient would be the singular cohomology with rational coefficient.

I would like to express my thanks to Professor Donu Arapura for suggesting me to look at this problem and for his encouragement. The definition of motivic dimension is due to him.

1. MAIN THEOREM

Let $V = Q_1 \cap \cdots \cap Q_k$ be a smooth complete intersection of k quadrics in a projective space $\mathbb{P}^{n+k}$. Then V is a smooth subvariety of $\mathbb{P}^{n+k}$ of $\dim V = n$. Consider the family of quadrics Q_t ($t \in \mathbb{P}^{k-1}$) with the base locus V . We can give more precise description of this family. Let $Q_l = \{F_l = 0\}$ for $l = 1, \dots, k$, where $F_l(x_0, x_1, \dots, x_{n+k}) = \sum_{i,j=0}^{n+k} c_{ij}^l x_i x_j$, $[c_{ij}^l]_{0 \leq i,j \leq n+k}$ is a symmetric $(n+k+1) \times (n+k+1)$ -matrix. Then for a general $t = (t_0, \dots, t_{k-1}) \in \mathbb{P}^{k-1}$, the fiber Q_t is given by the equation $\sum_{l=0}^{k-1} t_l F_{l+1}(x_0, \dots, x_{n+k}) = 0$. Let $\Delta \subset \mathbb{P}^{k-1}$ be the subvariety of $\mathbb{P}^{k-1}$ parametrizing all singular fibers in the family. Then Δ is a hypersurface in $\mathbb{P}^{k-1}$ of degree $n+k+1$. We assume that Δ is smooth. Set

$$X = \{(t, x) \in \mathbb{P}^{k-1} \times \mathbb{P}^{n+k} \mid x \in Q_t \subset \mathbb{P}^{n+k}\} \subset \mathbb{P}^{k-1} \times \mathbb{P}^{n+k}$$

and let $p_1 : X \rightarrow \mathbb{P}^{k-1}$ and $p_2 : X \rightarrow \mathbb{P}^{n+k}$ be projections. Note that X is a smooth projective variety of dimension $n+2k-2$.

$$(1) \quad \begin{array}{ccc} X & \xrightarrow{p_2} & \mathbb{P}^{n+k} \supset V \\ \downarrow p_1 & & \\ \mathbb{P}^{k-1} & & \end{array}$$

Let $i_X : X \hookrightarrow \mathbb{P}^{k-1} \times \mathbb{P}^{n+k}$ and $i_V : V \hookrightarrow \mathbb{P}^{n+k}$ be inclusions. Then by Lefschetz theorem, the restriction maps $i_X^* : H^{n+2k-2}(\mathbb{P}^{k-1} \times \mathbb{P}^{n+k}) \rightarrow H^{n+2k-2}(X)$ and $i_V^* : H^n(\mathbb{P}^{n+k}) \rightarrow H^n(V)$ are injections. Set

$$H_0^{n+2k-2}(X) = H^{n+2k-2}(X) / \text{im } i_X^*$$

and

$$H_0^n(V) = H^n(V) / \text{im } i_V^*$$

Remark 1.1. *In fact,*

$$H_0^{n+2k-2}(X) \cong \begin{cases} H_c^{n+2k-1}((\mathbb{P}^{k-1} \times \mathbb{P}^{n+k}) - X) & \text{if } n \text{ is even} \\ H^{n+2k-2}(X) & \text{if } n \text{ is odd} \end{cases}$$

and

$$H_0^n(V) \cong \begin{cases} H_c^{n+1}(\mathbb{P}^{n+k} - V) & \text{if } n \text{ is even} \\ H^n(V) & \text{if } n \text{ is odd} \end{cases}$$

as Hodge structures.

Now we can state our main theorem precisely:

Theorem 1.2. *Let V be a smooth complete intersection of k quadrics in $\mathbb{P}^{n+k}$. Then V has motivic dimension less than k . More precisely, there is a smooth projective variety W of dimension $k-1$ (resp. $k-2$) and surjection of rational Hodge structures*

$$\Theta : H^{k-1}(\tilde{W})(-q) \rightarrow H^n(V) \quad \text{if } n \text{ is odd and } k \text{ is even}$$

$$(\text{resp. } \Theta : H^{k-2}(\tilde{W})(-q) \rightarrow H^n(V) \quad \text{if } n \text{ is odd and } k \text{ is odd})$$

where $q = \frac{n-k+1}{2}$ (resp. $\frac{n-k+2}{2}$) and $\tilde{W}$ is a disjoint union of finitely many copies of W ,

$$\Theta : \bigoplus_r H^{2r}(W)(-q_r)^{\oplus l_r} \rightarrow H_0^n(V) \quad \text{if } n \text{ is even}$$

where $q_r = \frac{n-2r}{2}$ and l_r are positive integers given by explicit formula.

We will prove this theorem by considering two cases depending on the parity of $n + k$ in the last two sections. One immediate corollary of this theorem is

Corollary 1.3. *If $k \leq 4$, then Hodge conjecture holds for V .*

2. INTERMEDIATE STEP

Consider the projection $p_2 : X \rightarrow \mathbb{P}^{n+k}$. Note that for any $q = (q_0, \dots, q_{n+k}) \in \mathbb{P}^{n+k} - V$,

$$\begin{aligned} p_2^{-1}(q) &= \{(t, q) \in \mathbb{P}^{k-1} \times \mathbb{P}^{n+k} \mid q \in Q_t\} \\ &= \left\{ t = (t_0, \dots, t_{k-1}) \in \mathbb{P}^{k-1} \mid \sum_{l=0}^{k-1} \left(\sum_{i,j=0}^{n+k} c_{ij}^{l+1} q_i q_j \right) t_l = 0 \right\} \cong \mathbb{P}^{k-2} \end{aligned}$$

and for any $q \in V$,

$$p_2^{-1}(q) = \{(t, q) \in \mathbb{P}^{k-1} \times \mathbb{P}^{n+k} \mid q \in Q_t\} = \mathbb{P}^{k-1} \times \{q\}$$

since V is the base locus of the family. Hence we have the following diagram, which we will use throughout this section:

$$(2) \quad \begin{array}{ccccc} (\mathbb{P}^{n+k} - V) \times \mathbb{P}^{k-2} & \hookrightarrow & X & \xleftarrow{i_E} & \mathbb{P}^{k-1} \times V \\ \downarrow p_2 & & \downarrow p_2 & & \downarrow p_2 \\ \mathbb{P}^{n+k} - V & \hookrightarrow & \mathbb{P}^{n+k} & \xleftarrow{\quad} & V \end{array}$$

Lemma 2.1.

$$\dim H_0^n(V) = \dim H_0^{n+2k-2}(X)$$

Proof. From the diagram (2), we get

$$(3) \quad \chi(X) = (k-1)(n+k+1) + \chi(V)$$

Since X is a very ample divisor in $\mathbb{P}^{k-1} \times \mathbb{P}^{n+k}$ and V is a smooth complete intersection in $\mathbb{P}^{n+k}$, by Lefschetz Theorem we have

$$H^i(\mathbb{P}^{k-1} \times \mathbb{P}^{n+k}) \cong H^i(X), \quad H^j(\mathbb{P}^{n+k}) \cong H^j(V)$$

for any $i < \dim X = n + 2k - 2$ and $j < \dim V = n$.

(Case 1) If n is even : then we have

$$\chi(V) = 2 \sum_{i=0}^{n-1} (-1)^i b_i(V) + b_n(V) = n + b_n(V)$$

$$\begin{aligned} \chi(X) &= 2 \sum_{i=0}^{n+2k-3} (-1)^i b_i(X) + b_{n+2k-2}(X) \\ &= 2 \left(\frac{k(k+1)}{2} + k \cdot \frac{n-2}{2} \right) + b_{n+2k-2}(X) \\ &= nk + k^2 - k + b_{n+2k-2}(X) \end{aligned}$$

Therefore by (3) we get

$$nk + k^2 - k + b_{n+2k-2}(X) = (k-1)(n+k+1) + n + b_n(V)$$

i.e.

$$b_{n+2k-2}(X) = b_n(V) + k - 1$$

Note that

$$\dim H_0^{n+2k-2}(X) = b_{n+2k-2}(X) - \dim H^{n+2k-2}(\mathbb{P}^{k-1} \times \mathbb{P}^{n+k}) = b_{n+2k-2}(X) - k$$

and

$$\dim H_0^n(V) = b_n(V) - \dim H^n(\mathbb{P}^{n+k}) = b_n(V) - 1$$

therefore we get

$$\dim H_0^{n+2k-2}(X) = \dim H_0^n(V)$$

in this case.

(Case 2) If n is odd : then by the similar calculation, we get

$$\chi(X) = nk + k^2 - b_{n+2k-2}(X), \quad \chi(V) = (n+1) - b_n(V)$$

Then, again by (3)

$$nk + k^2 - b_{n+2k-2}(X) = (k-1)(n+k+1) + (n+1) - b_n(V) = nk + k^2 - b_n(V)$$

i.e.

$$b_{n+2k-2}(X) = b_n(V)$$

Hence, we get

$$\dim H_0^{n+2k-2}(X) = \dim H^{n+2k-2}(X) = \dim H^n(V) = \dim H_0^n(V)$$

in this case also, which finishes the proof of the Lemma. $\square$

Set $E = \mathbb{P}^{k-1} \times V \subset X$. Then we have a $\mathbb{P}^{k-1}$ -bundle $p_2 : E \rightarrow V$. Let $i_E : E \hookrightarrow X$ be an inclusion. Then we have the following of morphism of Hodge structures

$$\phi : H^n(V) \xrightarrow{p_2^*} H^n(E) \xrightarrow{i_{E*}} H^{n+2k-2}(X)$$

where $i_{E*} : H^n(E) \rightarrow H^{n+2k-2}(X)$ is the Gysin map. (Note that $\text{codim}(E, X) = k-1$)

Theorem 2.2. $\phi : H^n(V) \rightarrow H^{n+2k-2}(X)$ induces a morphism

$$\bar{\phi} : H_0^n(V)(-k+1) \rightarrow H_0^{n+2k-2}(X)$$

which is an isomorphism of rational Hodge structures.

Proof. First we prove that the induced morphism

$$\bar{\phi} : H_0^n(V) \rightarrow H_0^{n+2k-2}(X)$$

is well-defined. In the case when n is odd, then $\bar{\phi} = \phi$, hence the morphism is well-defined. In case when n is even, it is enough to show that $\phi(\text{im } i_V^*) \subseteq \text{im } i_X^*$. Let $[E] \in H^{2k-2}(X)$ be the fundamental class of E . Note that $i_X^* : H^{2k-2}(\mathbb{P}^{k-1} \times \mathbb{P}^{n+k}) \rightarrow H^{2k-2}(X)$ is an isomorphism by Lefschetz theorem since $2k-2 < \dim X = n+2k-2$. Hence there is $\gamma \in H^{2k-2}(\mathbb{P}^{k-1} \times \mathbb{P}^{n+k})$ such that $i_X^*(\gamma) = [E]$. In fact, we can write $\gamma = [\mathbb{P}^{k-1} \times S]$ where $[S] \in H^{2k-2}(\mathbb{P}^{n+k})$ such that

$$(4) \quad i_X^*(\gamma) = [X \cap (\mathbb{P}^{k-1} \times S)] = [\mathbb{P}^{k-1} \times V]$$

To show this, we consider the following commutative diagram:

$$\begin{array}{ccccc}
 H^0(E) & \xrightarrow{i_{*E}} & H^{2k-2}(X) & \xrightarrow{i_{X*}} & H^{2k}(\mathbb{P}^{k-1} \times \mathbb{P}^{n+k}) \\
 & \searrow^{(i_X \circ i_E)_*} & \uparrow i_X^* \cong & \nearrow \cong & \\
 & & H^{2k-2}(\mathbb{P}^{k-1} \times \mathbb{P}^{n+k}) & \xrightarrow{\cup[X]} &
 \end{array}$$

Since $\gamma \in H^{2k-2}(\mathbb{P}^{k-1} \times \mathbb{P}^{n+k})$, we may write $\gamma = \sum_{i=0}^{k-1} [a_i H^i \times b_i L^{k-1-i}]$, where H^i (resp. L^j) is a linear space in $\mathbb{P}^{k-1}$ (resp. $\mathbb{P}^{n+k}$) of codimension i (resp. j). By commutativity of the diagram, we have

$$[\mathbb{P}^{k-1} \times V] = i_{X*}[E] = i_{X*}i_X^*(\gamma) = \gamma \cup [X] = \sum_{i=0}^{k-1} [(a_i H^i \times b_i L^{k-1-i}) \cap X]$$

Hence $a_i = 0$ for $i \neq 0$, since $H^i \not\subseteq \mathbb{P}^{k-1}$ for $i \neq 0$ and then this forces $b_i = 0$ for $i \neq 0$ since $[b_i L^{k-1-i}] \notin H^{2k-2}(\mathbb{P}^{k-1} \times \mathbb{P}^{n+k})$ for $i > 0$. Hence

$$\gamma = [\mathbb{P}^{k-1} \times b_0 L^{k-1}]$$

We take S to be $b_0 L^{k-1}$ and we will use this S later.

Now consider the following diagram:

$$\begin{array}{ccccc}
 (5) & H^n(V) & \xrightarrow{p_2^*} & H^n(E) & \xrightarrow{i_{E*}} & H^{n+2k-2}(X) \\
 & \uparrow i_V^* & & \uparrow j_E^* & \swarrow i_E^* & \nearrow \cup[E] \\
 & & (I) & & (II) & H^n(X) & (IV) & & \uparrow i_X^* \\
 & & & & & \nearrow i_X^* & & & \\
 & H^n(\mathbb{P}^{n+k}) & \xrightarrow{pr_2^*} & H^n(\mathbb{P}^{k-1} \times \mathbb{P}^{n+k}) & \xrightarrow{\Phi} & H^{n+2k-2}(\mathbb{P}^{k-1} \times \mathbb{P}^{n+k})
 \end{array}$$

where

- (1) $j_E : E \rightarrow \mathbb{P}^{k-1} \times \mathbb{P}^{n+k}$, $i_E : E \rightarrow X$ and $i_X : X \rightarrow \mathbb{P}^{k-1} \times \mathbb{P}^{n+k}$ are inclusions,
- (2) $pr_2 : \mathbb{P}^{k-1} \times \mathbb{P}^{n+k} \rightarrow \mathbb{P}^{n+k}$ is the projection to the second factor,
- (3) $\Phi : H^n(\mathbb{P}^{k-1} \times \mathbb{P}^{n+k}) \rightarrow H^{n+2k-2}(\mathbb{P}^{k-1} \times \mathbb{P}^{n+k})$ is defined by $\Phi(\alpha) = \alpha \cup \gamma$.

Commutativity of (II) and (III) are clear and commutativity of (I) follows from the following commutative diagram:

$$\begin{array}{ccc}
 E = V \times \mathbb{P}^{n+k} & \xrightarrow{j_E} & \mathbb{P}^{k-1} \times \mathbb{P}^{n+k} \\
 \downarrow p_2 & & \downarrow pr_2 \\
 V & \xrightarrow{i_V} & \mathbb{P}^{n+k}
 \end{array}$$

We show the commutativity of (IV). Let $\alpha \in H^n(\mathbb{P}^{k-1} \times \mathbb{P}^{n+k})$. Then

$$i_X^* \circ \Phi(\alpha) = i_X^*(\alpha \cup \gamma) = i_X^*(\alpha) \cup i_X^*(\gamma) = i_X^*(\alpha) \cup [E]$$

by the definition of γ . Hence (IV) commutes. Therefore we have

$$\phi \circ i_V^* = i_X^* \circ \Phi \circ pr_2^*$$

and hence $\phi(\text{im } i_V^*) \subseteq \text{im } i_X^*$ and ϕ induces a well-defined morphism $\bar{\phi} : H_0^n(V) \rightarrow H_0^{n+2k-2}(X)$ in this case also.

To show that $\bar{\phi}$ is an isomorphism, note that we have

$$(6) \quad H^n(E) \cong (H^n(V) \otimes H^0(\mathbb{P}^{k-1})) \oplus \left(\bigoplus_{i=1}^{k-1} H^{n-2i}(\mathbb{P}^{n+k}) \otimes H^{2i}(\mathbb{P}^{k-1}) \right)$$

by the Künneth formula and Lefschetz theorem.

(Case 1) If n is odd : in this case $\phi = \bar{\phi}$ and we show that $\phi = i_{E*} \circ p_2^*$ is surjective. Since n is odd, $H^{n-2i}(\mathbb{P}^{k-1}) = 0$ for all i . Hence (6) gives an isomorphism

$$p_2^* : H^n(V) \xrightarrow{\cong} H^n(E)$$

Now for the morphism $i_{E*} : H^n(E) \rightarrow H^{n+2k-2}(X)$, consider the following Gysin exact sequence

$$(7) \quad \cdots \rightarrow H^{n+2k-3}(X-E) \rightarrow H^n(E) \xrightarrow{i_{E*}} H^{n+2k-2}(X) \rightarrow H^{n+2k-2}(X-E) \rightarrow \cdots$$

Since $X - E \cong \mathbb{P}^{k-2} \times (\mathbb{P}^{n+k} - V)$,

$$\begin{aligned} H^{n+2k-2}(X-E) &= H^{n+2k-2}(\mathbb{P}^{k-2} \times (\mathbb{P}^{n+k} - V)) \\ &= \bigoplus_{i=0}^{k-2} H^{(n+2k-2)-2i}(\mathbb{P}^{n+k} - V)(-i) = 0 \end{aligned}$$

Hence, $i_{E*} : H^n(E) \rightarrow H^{n+2k-2}(X)$ is surjective and hence we get the surjection $\phi = i_{E*} \circ p_2^*$. Now lemma 2.1 implies that $\bar{\phi} = \phi$ is an isomorphism in this case.

(Case 2) If $n = 2l$ is even : then (6) gives that $p_2^* : H^n(V) \rightarrow H^n(E)$ is an injection. Since $X - E \cong \mathbb{P}^{k-2} \times (\mathbb{P}^{n+k} - V)$, we get

$$H^{n+2k-3}(E) = H^{n+2k-3}(\mathbb{P}^{k-2} \times (\mathbb{P}^{n+k} - V)) = \bigoplus_{i=0}^{k-2} H^{(n+2k-3)-2i}(\mathbb{P}^{n+k} - V)(-i) = 0$$

Then Gysin exact sequence (7) implies that $i_{E*} : H^n(E) \rightarrow H^{n+2k-2}(X)$ is injective. Since $\phi = i_{E*} \circ p_2^*$, we get the injection $\phi = i_{E*} \circ p_2^* : H^n(V) \rightarrow H^{n+2k-2}(X)$. We show that the induced map $\bar{\phi}$ is also an injection in this case: Suppose $\bar{\phi}(\bar{\alpha}) = 0$. Then

$$\phi(\alpha) \in \text{im}[i_X^* : H^{n+2k-2}(\mathbb{P}^{k-1} \times \mathbb{P}^{n+k}) \rightarrow H^{n+2k-2}(X)]$$

where $\alpha \in H^n(V)$ which maps to $\bar{\alpha} \in H_0^n(V)$. Let $\beta \in H^{n+2k-2}(\mathbb{P}^{k-1} \times \mathbb{P}^{n+k})$ such that $i_X^*(\beta) = \phi(\alpha)$. We claim that β is a cycle in $H^0(\mathbb{P}^{k-1}) \otimes H^{n+2k-2}(\mathbb{P}^{n+k})$.

$$\begin{array}{ccccc} & & \xrightarrow{\cup c_{k-1}(N_{E/X})} & & H^{n+2k-2}(E) \\ & & \nearrow i_{E*} & & \nearrow i_E^* \\ H^n(V) & \xrightarrow{p_2^*} & H^n(E) & \xrightarrow{i_{E*}} & H^{n+2k-2}(X) \\ \uparrow i_V^* & & \uparrow & & \uparrow i_X^* \\ H^n(\mathbb{P}^{n+k}) & \xrightarrow{pr_2^*} & H^n(\mathbb{P}^{k-1} \times \mathbb{P}^{n+k}) & \xrightarrow{\phi} & H^{n+2k-2}(\mathbb{P}^{k-1} \times \mathbb{P}^{n+k}) \end{array}$$

$(i_X \circ i_E)^*$

By applying i_E^* to $\phi(\alpha) = i_X^*(\beta) \in H^{n+2k-2}(X)$, we get

$$(8) \quad i_E^*(\phi(\alpha)) = i_E^* i_{E*} p_2^*(\alpha) = p_2^*(\alpha) \cup c_{k-1}(N_{E/X}) = i_E^*(i_X^*(\beta)) = (i_X \circ i_E)^*(\beta)$$

where the second equality comes from the self-intersection formula ([F, p103]).

From the inclusions $E \xrightarrow{i_E} X \xrightarrow{i_X} \mathbb{P}^{k-1} \times \mathbb{P}^{n+k}$, we have

$$0 \rightarrow i_E^*(N_{X/\mathbb{P}^{k-1} \times \mathbb{P}^{n+k}}) \rightarrow N_{E/\mathbb{P}^{k-1} \times \mathbb{P}^{n+k}} \rightarrow N_{E/X} \rightarrow 0$$

Hence

$$c(N_{E/\mathbb{P}^{k-1} \times \mathbb{P}^{n+k}}) = c(i_E^*(N_{X/\mathbb{P}^{k-1} \times \mathbb{P}^{n+k}})) \cdot c(N_{E/X})$$

In particular,

$$(9) \quad c_k(N_{E/\mathbb{P}^{k-1} \times \mathbb{P}^{n+k}}) = c_1(i_E^*(N_{X/\mathbb{P}^{k-1} \times \mathbb{P}^{n+k}})) \cdot c_{k-1}(N_{E/X}) = i_E^*([X]|_X) \cup_E [E]|_E$$

since X is a divisor in $\mathbb{P}^{k-1} \times \mathbb{P}^{n+k}$, where $\cup_E$ is the cup product on E . Since the inclusion $i_X \circ i_E : E \rightarrow \mathbb{P}^{k-1} \times \mathbb{P}^{n+k}$ is actually $(\text{id}_{\mathbb{P}^{k-1}}, i_V) : \mathbb{P}^{k-1} \times V \hookrightarrow \mathbb{P}^{k-1} \times \mathbb{P}^{n+k}$, we have $N_{E/\mathbb{P}^{k-1} \times \mathbb{P}^{n+k}} = p_2^*(N_{V/\mathbb{P}^{n+k}})$ and hence

$$(10) \quad c_k(N_{E/\mathbb{P}^{k-1} \times \mathbb{P}^{n+k}}) = p_2^*(c_k(N_{V/\mathbb{P}^{n+k}})) \in H^0(\mathbb{P}^{k-1}) \otimes H^{2k}(V) \subset H^{2k}(E)$$

Now since X is an ample divisor in $\mathbb{P}^{k-1} \times \mathbb{P}^{n+k}$, $[X] = [aH + bL] \in H^2(\mathbb{P}^{k-1} \times \mathbb{P}^{n+k})$ where H (resp. L) is a hyperplane in $\mathbb{P}^{k-1}$ (resp. $\mathbb{P}^{n+k}$) and $a, b \geq 0$ such that $ab \neq 0$. Now by using (4),

$$\begin{aligned} i_E^*([X]|_X) \cup_E [E]|_E &= i_E^*([X]|_X \cup_X [E]) = i_E^*([X]|_X \cup_X i_X^*(\gamma)) \\ &= i_E^* i_X^*([X] \cup_{\mathbb{P}^{k-1} \times \mathbb{P}^{n+k}} \gamma) \\ &= i_E^* i_X^*([aH + bL] \cup_{\mathbb{P}^{k-1} \times \mathbb{P}^{n+k}} [\mathbb{P}^{k-1} \times S]) \\ &= i_E^* i_X^*([a(H \times S) + b(\mathbb{P}^{k-1} \times (S \cap L))]) \\ &\in (H^2(\mathbb{P}^{k-1}) \otimes H^{2k-2}(V)) \oplus (H^0(\mathbb{P}^{k-1}) \otimes H^{2k}(V)) \end{aligned}$$

Then by (9), (10), we have $a = 0$ and hence $[X] = [bL] \in H^2(\mathbb{P}^{k-1} \times \mathbb{P}^{n+k})$. Set P be a hypersurface of degree b in $\mathbb{P}^{n+k}$ such that $[X] = [\mathbb{P}^{k-1} \times P]$. Since X contains $E = \mathbb{P}^{k-1} \times V$, we may assume that $V \subset P$. Then $N_{E/X} = p_2^*(N_{V/P})$, and hence

$$c_{k-1}(N_{E/X}) = p_2^*(c_{k-1}(N_{V/P})) \in H^0(\mathbb{P}^{k-1}) \otimes H^{2k-2}(V)$$

Now, since $p_2^*(\alpha) \in H^0(\mathbb{P}^{k-1}) \otimes H^n(V)$, we have

$$i_E^* \phi(\alpha) = p_2^*(\alpha) \cup c_{k-1}(N_{E/X}) \in H^0(\mathbb{P}^{k-1}) \otimes H^{n+2k-2}(V)$$

Then by (8), we have

$$i_E^* \phi(\alpha) = (i_E \circ i_X)^*(\beta) = (\text{id}_{\mathbb{P}^{k-1}}, i_V^*)(\beta) \in H^0(\mathbb{P}^{k-1}) \otimes H^{n+2k-2}(V)$$

Hence $\beta \in H^0(\mathbb{P}^{k-1}) \otimes H^{n+2k-2}(\mathbb{P}^{n+k})$.

Next note that by cupping with $\gamma = [\mathbb{P}^{k-1} \times S]$, $\Phi : H^n(\mathbb{P}^{k-1} \times \mathbb{P}^{n+k}) \rightarrow H^{n+2k-2}(\mathbb{P}^{k-1} \times \mathbb{P}^{n+k})$ maps the Künneth component $H^i(\mathbb{P}^{k-1}) \otimes H^{n-i}(\mathbb{P}^{n+k})$ of $H^n(\mathbb{P}^{k-1} \times \mathbb{P}^{n+k})$ to the Künneth component $H^i(\mathbb{P}^{k-1}) \otimes H^{n+2k-2-i}(\mathbb{P}^{n+k})$ of $H^{n+2k-2}(\mathbb{P}^{k-1} \times \mathbb{P}^{n+k})$ isomorphically. Hence we have $\eta \in H^0(\mathbb{P}^{k-1}) \otimes H^n(\mathbb{P}^{n+k})$ such that $\eta \cup \gamma = \Phi(\eta) = \beta$. Now $\eta \in \text{im}[pr_2^* : H^n(\mathbb{P}^{n+k}) \rightarrow H^n(\mathbb{P}^{k-1} \times \mathbb{P}^{n+k})]$ and we may consider $\eta \in H^n(\mathbb{P}^{n+k})$ since pr_2^* is injective. Then by commutativity of diagram (5), we have

$$\phi \circ i_V^*(\eta) = i_X^* \circ \Phi \circ pr_2^*(\eta) = i_X^*(\eta \cup \gamma) = i_X^*(\beta) = \phi(\alpha)$$

Since ϕ is injective, we have $\alpha = i_V^*(\eta)$. Hence $\bar{\phi}$ is also an injection. Now by lemma 2.1, we can conclude that $\bar{\phi} : H_0^n(V) \rightarrow H_0^{n+2k-2}(X)$ is an isomorphism. $\square$

3. PROOF OF THEOREM 1.2 WHEN $n + k$ IS ODD

Throughout this section, we assume that $n + k = 2m + 1$.

In order to prove theorem 1.2, we use the construction of O'Grady[O]. We give a brief outline of his construction here. For detailed construction, see [O].

Recall the diagram (1) and consider the projection $p_1 : X \rightarrow \mathbb{P}^{k-1}$ and recall that Δ is the discriminant variety, which is a smooth hypersurface in $\mathbb{P}^{k-1}$ by our assumption. In case when $n + k$ is odd, for a general $t \in \mathbb{P}^{k-1}$, the fiber $p_1^{-1}(t) = Q_t$ is a smooth quadric of dimension $n + k - 1 = 2m$ in $\mathbb{P}^{n+k}$. Hence it contains two irreducible families of m -planes parametrized by F_t^1 and F_t^2 . Note that $F_t^1 \cong F_t^2$ and $\dim F_t^i = \frac{m(m+1)}{2}$ [GH]. Let F be the abstract variety to which F_t^i is isomorphic for $i = 1, 2$ and for any t . Let W be a double covering of $\mathbb{P}^{k-1}$ branched over Δ and let $\sigma : W \rightarrow \mathbb{P}^{k-1}$ be the covering map. Set

$$P = \{M \subset X \mid p_1(M) = t \in \mathbb{P}^{k-1} \text{ a point}, p_2(M) \cong \mathbb{P}^m \subset \mathbb{P}^{n+k}\}$$

Then there is a natural map $\psi : P \rightarrow \mathbb{P}^{k-1}$ defined by $\psi(M) = p_1(M) \in \mathbb{P}^{k-1}$. Then the Stein factorization of ψ is factored through W and get a composition

$$\psi : P \xrightarrow{f} W \xrightarrow{\sigma} \mathbb{P}^{k-1}$$

and set

$$\begin{aligned} \Gamma &= \{(M, x) \in P \times X \mid x \in M \subset X\} \\ &= \{(M, x) \in P \times X \mid p_1(x) = p_1(M) \in \mathbb{P}^{k-1}, p_2(x) \in p_2(M) \cong \mathbb{P}^m\} \end{aligned}$$

Let $pr_1 : \Gamma \rightarrow P$ and $pr_2 : \Gamma \rightarrow X$ be the projections. We summarize the construction in the following diagram:

$$\begin{array}{ccccc} & & \Gamma & & \\ & \swarrow pr_1 & & \searrow pr_2 & \\ P & & & & X \\ & \searrow \psi & & \swarrow p_1 & \\ W & \xrightarrow{\sigma} & & & \mathbb{P}^{k-1} \end{array}$$

Note that for any $w \in W$,

$$f^{-1}(w) = \{M \in P \mid p_1(M) = \sigma(w), p_2(M) \cong \mathbb{P}^m \in \mathbb{P}^{n+k}\} \cong \{\mathbb{P}^m \subset Q_{\sigma(w)}\} \cong F$$

and for any $M \in P$,

$$\begin{aligned} pr_1^{-1}(M) &= \{(M, x) \in P \times X \mid x \in M \subset X\} \\ &\cong \{x \in X \mid p_1(x) = p_1(M), p_2(x) \in p_2(M) \cong \mathbb{P}^m\} \cong \mathbb{P}^m \end{aligned}$$

Hence $f : P \rightarrow W$ is F -bundle and $pr_1 : \Gamma \rightarrow P$ is $\mathbb{P}^m$ -bundle.

Now we give a proof of theorem 1.2 in case of $n + k$ odd.

Proof of theorem 1.2 when $n + k$ is odd. Consider the morphism of Hodge structures

$$pr_{2*} : H_{n+2k-2}(\Gamma) \rightarrow H_{n+2k-2}(X)$$

We show that pr_{2*} is a surjection. First note that $pr_2 : \Gamma \rightarrow X$ is a surjection. Hence we can take an iterated hyperplane section Γ_1 of Γ such that $\dim \Gamma_1 = \dim X = n + 2k - 2$ and Γ_1 surjects onto X . Let $g = pr_2|_{\Gamma_1} : \Gamma_1 \rightarrow X$. Then g is a generically finite map. Let $j : \Gamma_1 \rightarrow \Gamma$ be an inclusion. Then we have the following commutative diagram:

where $l_j = \dim H_{n+k-1-2j}(F)$. Therefore we have

$$\begin{aligned} H_{n+2k-2}(\Gamma) &= \bigoplus_{j=0}^m H_{n+2k-2-2j}(P)(j) \\ &\cong \bigoplus_{j=0}^m (H_{k-1}(W)(m-j)^{\oplus l_j})(j) = H_{k-1}(W)(m)^{\oplus N} \end{aligned}$$

where $N = \sum_{j=0}^m l_j$. Thus, we have a surjection

$$pr_{2*} : H_{k-1}(W)(m)^{\oplus N} \rightarrow H_{n+2k-2}(X)$$

and by Poincaré duality, we get a surjection

$$H^{k-1}(W)(-m)^{\oplus N} \longrightarrow H^{n+2k-2}(X)$$

By choosing $\tilde{W}$ to be the disjoint union of N copies of W and by composing with $\bar{\phi}^{-1}$ in theorem 2.2, we get a surjection

$$\Theta : H^{k-1}(\tilde{W})(-m+k-1) \longrightarrow H^n(V)$$

as we claimed.

(Case 2) If n is even (and hence k is odd) : in this case, (11) gives

$$H^{2d-(n+2k-2-2j)}(P) \cong \bigoplus_r (H^{2r}(W) \otimes H^{2d-(n+2k-2-2j)-2r}(F))$$

for each j , or equivalently

$$H_{n+2k-2-2j}(P) \cong \bigoplus_s (H_{2s}(W) \otimes H_{n+2k-2-2j-2s}(F))$$

Therefore we have

$$\begin{aligned} H_{n+2k-2}(\Gamma) &= \bigoplus_{j=0}^m H_{n+2k-2-2j}(P)(j) \\ &\cong \bigoplus_{j=0}^m \bigoplus_s (H_{2s}(W) \otimes H_{n+2k-2-2j-2s}(F))(j) \\ &= \bigoplus_s H_{2s}(W) \otimes \left(\bigoplus_{j=0}^m H_{n+2k-2-2j-2s}(F) \right)(j) \\ &= \bigoplus_s H_{2s}(W)(q_s)^{\oplus l_s} \end{aligned}$$

where $q_s = \frac{n+2(k-1)-2s}{2}$ and $l_s = \sum_{j=0}^m \dim H_{n+2k-2-2j-2s}(F)$. Thus, we have a surjection

$$pr_{2*} : \bigoplus_s H_{2s}(W)(q_s)^{\oplus l_s} \longrightarrow H_{n+2k-2}(X)$$

Recall that $H_0^{n+2k-2}(X) = H^{n+2k-2}(X)/\text{im } i_X^*$. Hence we get a composition of surjections

$$\bigoplus_s H_{2s}(W)(-q_s)^{\oplus l_s} \longrightarrow H^{n+2k-2}(X) \longrightarrow H_0^{n+2k-2}(X)$$

Therefore, by composing with $\bar{\phi}^{-1}$ from theorem 2.2 we get a surjection of Hodge structures

$$\Theta : \bigoplus_s H^{2s}(W)(-q_s + k - 1)^{\oplus l_s} \longrightarrow H_0^n(V)$$

(Note that $q_s - k + 1 = \frac{n-2s}{2}$.) This completes the proof of theorem 1.2 in case when $n + k$ is odd. $\square$

4. PROOF OF THEOREM 1.2 WHEN $n + k$ IS EVEN

Throughout this section, we assume that $n + k = 2m$ is even.

Again we start by considering the projection $p_1 : X \rightarrow \mathbb{P}^{k-1}$ in diagram (1). Recall that the discriminant variety Δ is a smooth hypersurface of degree $n + k + 1$ in $\mathbb{P}^{k-1}$ by our assumption. Set $U = \mathbb{P}^{k-1} - \Delta$ and let $X_\Delta = p_1^{-1}(\Delta)$ and $X_U = p_1^{-1}(U)$.

$$(12) \quad \begin{array}{ccccc} X_\Delta & \hookrightarrow & X & \longleftarrow & X_U \\ \downarrow p_1 & & \downarrow p_1 & & \downarrow p_1 \\ \Delta & \hookrightarrow & \mathbb{P}^{k-1} & \longleftarrow & U = \mathbb{P}^{k-1} - \Delta \end{array}$$

Note that for any $t \in \Delta$, $p_1^{-1}(t)$ is a singular quadric in the family. Since we have assumed that Δ is smooth, a singular fiber is a cone through a 0-plane (i.e. a point) over a quadric of rank $n + k$ in $\mathbb{P}^{n+k-1}$, i.e. all singular fibers are cones over a smooth quadric $\tilde{Q}_t$ of dimension $n + k - 2$ in $\mathbb{P}^{n+k-1}$. For any $t \in \Delta$, we denote $p_1^{-1}(t) = C_t$ a cone with a vertex 0_t . We can form a family of quadrics of dimension $n + k - 2$ over Δ as follows: Let $s : \Delta \rightarrow X_\Delta$ be a section of p_1 defined by $s(t) = 0_t$ for any $t \in \Delta$. Let Y_0 be a general hyperplane section of $X_\Delta - s(\Delta)$ and let $p_1|_{Y_0} : Y_0 \rightarrow \Delta$ be the obvious map. Let Y be a smooth compactification of Y_0 . Then by using theorem by Hironaka, we may assume that the rational map $Y \rightarrow \Delta$ is actually a morphism. We denote this morphism by $\pi : Y \rightarrow \Delta$. Then for a general $t \in \Delta$, $\pi^{-1}(t) = \tilde{Q}_t$ a smooth quadric of dimension $n + k - 2$ in $\mathbb{P}^{n+k-1}$.

$$\begin{array}{ccccccc} Y & \longleftarrow & Y_0 & \hookrightarrow & X_\Delta & \hookrightarrow & X \\ & \searrow \pi & \downarrow p_1|_{Y_0} & & \downarrow p_1 & & \downarrow p_1 \\ & & \Delta & \xlongequal{\quad} & \Delta & \hookrightarrow & \mathbb{P}^{k-1} \end{array}$$

Lemma 4.1. *For any p*

$$H^p(X_\Delta) \cong H^{p-2}(Y)(-1) \quad \text{as Hodge structures}$$

Proof. Consider the Leray spectral sequence associated to the map $p_1 : X_\Delta \rightarrow \Delta$

$${}^I E_2^{pq} = H^p(\Delta, R^q p_{1*} \mathbb{Q}) \Rightarrow H^{p+q}(X_\Delta, \mathbb{Q})$$

and one associated to the map $\pi : Y \rightarrow \Delta$

$${}^{II} E_2^{pq} = H^p(\Delta, R^q \pi_* \mathbb{Q}) \Rightarrow H^{p+q}(Y, \mathbb{Q})$$

Since for any $t \in \Delta$, C_t is a cone through a point 0_t over a smooth quadric $\tilde{Q}_t$ in $\mathbb{P}^{n+k-1}$, we have

$$\begin{aligned} R^q p_{1*} \mathbb{Q} &= R^q (p_1|_{Y_0})_* \mathbb{Q} = (R^{2(n+k-1)-q} (p_1|_{Y_0})_* \mathbb{Q})^*(-n-k+1) \\ &= (R^{2(n+k-1)-q} \pi_* \mathbb{Q})^*(-n-k+1) = R^{q-2} \pi_* \mathbb{Q}(-1) \end{aligned}$$

Hence we have

$$'E_2^{pq} = ''E_2^{p,q-2}(-1)$$

and so

$$H^{p+q}(X_\Delta) \cong H^{p+q-2}(Y)(-1)$$

as Hodge structures [A]. $\square$

In particular,

$$(13) \quad H^{n+2k-2}(X_\Delta) \cong H^{n+2k-4}(Y)(-1)$$

Lemma 4.2. *There is an injection of Hodge structures*

$$0 \rightarrow H_0^{n+2k-2}(X) \rightarrow H^{n+2k-4}(Y)(-1)$$

Proof. From the top row of the diagram (12), we have an exact sequence of mixed Hodge structures

$$(14) \quad \dots \rightarrow H_c^{n+2k-2}(X_U) \rightarrow H^{n+2k-2}(X) \rightarrow H^{n+2k-2}(X_\Delta) \rightarrow \dots$$

By using (13) and by taking the exact functor Gr_{n+2k-2}^W , we get

$$(15) \quad 0 \rightarrow Gr_{n+2k-2}^W H_c^{n+2k-2}(X_U) \rightarrow H^{n+2k-2}(X) \rightarrow H^{n+2k-4}(Y)(-1) \rightarrow \dots$$

Consider the morphism $p_1 : X_U \rightarrow U$ and the Leray spectral sequence associated to it

$$E_2^{pq} = H_c^p(U, R^q p_{1*} \mathbb{Q}) \Rightarrow H_c^{p+q}(X_U, \mathbb{Q})$$

Note that

$$E_2^{pq} = \begin{cases} 0 & \text{if } q \text{ is odd} \\ H_c^p(U, \mathbb{Q}) & \text{if } q \text{ is even} \end{cases}$$

since $(R^q p_{1*} \mathbb{Q})_t = H^q(Q_t, \mathbb{Q})$ and Q_t is a smooth quadric of dimension $n+k-1$ (odd). Also, we have

$$0 \rightarrow H_c^0(U) \rightarrow H^0(\mathbb{P}^{k-1}) \rightarrow H^0(\Delta) \rightarrow H_c^1(U) \rightarrow 0$$

$$0 \rightarrow H^{2i-1}(\Delta) \rightarrow H_c^{2i}(U) \rightarrow H^{2i}(\mathbb{P}^{k-1}) \rightarrow H^{2i}(\Delta) \rightarrow H_c^{2i+1}(U) \rightarrow 0$$

for $1 \leq i \leq k-2$ and

$$H_c^{2k-2}(U) \cong H^{2k-2}(\mathbb{P}^{k-1})$$

Now, since Δ is a smooth hypersurface in $\mathbb{P}^{k-1}$, by Lefschetz theorem we get

$$H^j(\mathbb{P}^{k-1}) \xrightarrow{\cong} H^j(\Delta), \quad \text{for } j < \dim \Delta = k-2$$

Hence $H^{2i-1}(\Delta) = 0$ and hence $H_c^{2i}(U) = H_c^{2i+1}(U) = 0$ for i such that $2i < \dim \Delta = k-2$. By applying duality on $H^{2i-1}(\Delta)$, we get $H_c^j(U) = 0$ unless $j = k-1$ or $j = 2k-2$. Hence the Leray spectral sequence degenerates at E_2 and we have a short exact sequence

$$0 \rightarrow E_\infty^{2k-2,n} \rightarrow H_c^{n+2k-2}(X_U) \rightarrow E_\infty^{k-1,n+k-1} \rightarrow 0$$

Note that $E_\infty^{k-1,n+k-1} = E_2^{k-1,n+k-1} = 0$ since $n+k$ is even. Hence we have

$$(16) \quad E_\infty^{2k-2,n} \cong H_c^{n+2k-2}(X_U)$$

(Case 1) If n is odd : Then $E_\infty^{2k-2,n} = 0$ also and hence we have $H_c^{n+2k-2}(X_U) = 0$. So (15) gives an injection

$$0 \rightarrow H^{n+2k-2}(X) \rightarrow H^{n+2k-4}(Y)(-1)$$

(Case 2) If n is even : In this case, (16) gives

$$\begin{aligned} H_c^{n+2k-2}(X_U) &\cong E_\infty^{2k-2,n} = E_2^{2k-2,n} = H_c^{2k-2}(U, R^n p_{1*} \mathbb{Q}) \\ &\cong H_c^{2k-2}(U) \otimes H^n(Q_t) \cong H^{2k-2}(\mathbb{P}^{k-1}) \otimes H^n(\mathbb{P}^{n+k}) \end{aligned}$$

by Lefschetz theorem. Hence, we can rewrite the exact sequence (15) as follows:

$$0 \rightarrow H^{2k-2}(\mathbb{P}^{k-1}) \otimes H^n(\mathbb{P}^{n+k}) \xrightarrow{h_*} H^{n+2k-2}(X) \rightarrow H^{n+2k-4}(Y) \rightarrow \dots$$

Recall that $H_0^{n+2k-2}(X) \cong H^{n+2k-2}(X)/\text{im } i_X^*$ and note that $\text{im } h_* \cap H_0^{n+2k-2}(X) = \emptyset$: In fact, h_* can be factorized as

$$H^{2k-2}(\mathbb{P}^{k-1}) \otimes H^n(\mathbb{P}^{n+k}) \xrightarrow{\quad \quad \quad h_* \quad \quad \quad} H^{n+2k-2}(\mathbb{P}^{k-1} \times \mathbb{P}^{n+k}) \xrightarrow{i_X^*} H^{n+2k-2}(X)$$

Hence $\text{im } h_* \subseteq \text{im } i_X^*$, and we get an injection

$$0 \rightarrow H_0^{n+2k-2}(X) \rightarrow H^{n+2k-4}(Y)(-1)$$

□

Now we have a family $\pi : Y \rightarrow \Delta$ of quadrics of dimension $n+k-2$ which is even, so we can form a double covering of Δ as in the construction of O'Grady([O]). For a general $t \in \Delta$, the fiber $\tilde{Q}_t$, which is a smooth quadric of dimension $n+k-2 = 2(m-1)$, contains two $\frac{m(m-1)}{2}$ -dimensional irreducible families F_t^1, F_t^2 of $(m-1)$ -planes[GH]. As in the case of $n+k$ odd, we can form the following diagram

$$\begin{array}{ccccc} & & \Gamma & & \\ & \swarrow pr_1 & & \searrow pr_2 & \\ P & & & & Y \xrightarrow{\pi_2} \mathbb{P}^{n+k-1} \\ & \searrow \psi & & \downarrow \pi & \\ W & \xrightarrow{\sigma} & \Delta & & \end{array}$$

where

- (1) $P = \{M \subset Y \mid \pi_1(M) = t \in \Delta \text{ a point}, \pi_2(M) \cong \mathbb{P}^{m-1} \subset \mathbb{P}^{n+k-1}\}$
- (2) $\Gamma = \{(M, y) \in P \times Y \mid \pi(y) = \pi(M) \in \Delta, \pi(y) \in \pi_2(M) \cong \mathbb{P}^{m-1} \subset \tilde{Q}_{\pi(M)}\}$, pr_1 and pr_2 are projections.
- (3) $\psi : P \rightarrow \Delta$ is a natural map defined by $\psi(M) = \pi(M) = t \in \Delta$
- (4) $f : P \rightarrow W$ is F -bundle and $pr_1 : \Gamma \rightarrow P$ is $\mathbb{P}^{m-1}$ -bundle, where F is the abstract variety such that F_t^i is isomorphic to it for $i = 1, 2$ and $t \in \Delta$
- (5) $\sigma : W \rightarrow \Delta$ is a double covering branched over the discriminant variety Δ_1 of the family $\pi : Y \rightarrow \Delta$.

Lemma 4.3. *There are surjections of Hodge structures*

(1)

$$H^{k-2}(\tilde{W})(-m+1) \rightarrow H^{n+2k-4}(Y) \quad \text{if } n \text{ is odd}$$

where $\tilde{W}$ is a disjoint union of finitely many copies of W ,

(2)

$$\bigoplus_r H^{2r}(W)(-q_r)^{\oplus l_r} \rightarrow H^{n+2k-4}(Y) \quad \text{if } n \text{ is even}$$

where $q_r = \frac{n+2k-4-2r}{2}$ and $l_r = \sum_j \dim H_{n+2k-4-2j-2r}(F)$.

Proof. First note that by using the same arguments in the proof of theorem 1.2 in the case of $n + k$ odd, we can show that

$$(17) \quad pr_{2*} : H_{n+2k-4}(\Gamma) \rightarrow H_{n+2k-4}(Y)$$

is surjective and

$$\begin{aligned} H_{n+2k-4}(\Gamma) &= \bigoplus_{j=0}^{m-1} (H_{n+2k-4-2j}(P) \otimes H_{2j}(\mathbb{P}^{m-1})) \\ &= \bigoplus_{j=0}^{m-1} H_{n+2k-4-2j}(P)(j) \\ &= \bigoplus_{j=0}^{m-1} \bigoplus_s (H_{n+2k-4-2j-2s}(W) \otimes H_{2s}(F))(j) \end{aligned}$$

(Case 1) If n is odd : In this case, $n + 2k - 4 - 2j - 2s$ is odd. But $H_r(W)$ is zero for an odd number r unless $r = k - 2 = \dim W$. Hence we have

$$H_{n+2k-4-2j}(P) = H_{k-2}(W) \otimes H_{n+k-2-2j}(F) = H_{k-2}(W)(m-1-j)^{\oplus l_j}$$

where $l_j = \dim H_{n+k-2-2j}(F)$. Thus we have

$$\begin{aligned} H_{n+2k-4}(\Gamma) &= \bigoplus_{j=0}^{m-1} H_{k-2}(W)(m-1-j)^{\oplus l_j}(j) \\ &= \bigoplus_{j=0}^{m-1} H_{k-2}(W)(m-1)^{\oplus l_j} = H_{k-2}(W)(m-1)^{\oplus N} \end{aligned}$$

where $N = \sum_{j=0}^m l_j$. Hence by (17) and using Poincaré duality, we get a surjection

$$pr_{2*} : H^{k-2}(W)(-m+1)^{\oplus N} \longrightarrow H^{n+k-4}(Y)$$

By setting $\tilde{W}$ to be a disjoint union of N copies of W , we get the desired surjection in this case.

(Case 2) If n is even : In this case, $n + 2k - 4 - 2j$ is even. Hence we have

$$H_{n+2k-4-2j}(P) = \bigoplus_s (H_{2s}(W) \otimes H_{n+2k-4-2j-2s}(F))$$

Thus we have

$$\begin{aligned} H_{n+2k-4}(\Gamma) &= \bigoplus_{j=0}^{m-1} H_{n+2k-4-2j}(P)(j) \\ &= \bigoplus_{j=0}^{m-1} \left(\bigoplus_s (H_{2s}(W) \otimes H_{n+2k-4-2j-2s}(F)) \right) (j) \\ &= \bigoplus_s H_{2s}(W) \otimes \left(\bigoplus_{j=0}^{m-1} H_{n+2k-4-2j-2s}(F) \right) (j) \\ &= \bigoplus_s H_{2s}(W)(q_s)^{\oplus l_s} \end{aligned}$$

where $q_s = \frac{n+2k-4-2s}{2}$ and $l_s = \sum_{j=0}^{m-1} \dim H_{n+2k-4-2j-2s}(F)$. Hence by (17) and using Poincaré duality, we get a surjection

$$pr_{2*} : \bigoplus_r H^{2r}(W)(-q_r)^{\oplus l_r} \twoheadrightarrow H^{n+2k-4}(Y)$$

in this case. □

We can summarize lemmas 4.2 and 4.3 in the following diagram:

(18)

$$\begin{array}{ccccccc} & & H^*(W) & & & & \\ & & \downarrow \theta & \searrow \psi_2 \circ \theta & & & \\ 0 \rightarrow & H_0^{n+2k-2}(X) & \xrightarrow{\psi_1} & H^{n+2k-4}(Y)(-1) & \xrightarrow{\psi_2} & Gr_{n+2k-2}^W H_c^{n+2k-1}(X_U) & \rightarrow 0 \end{array}$$

where with the notation in theorem 4.3

$$H^*(W) = \begin{cases} H^{k-2}(\tilde{W})(-m) & \text{if } n \text{ is odd} \\ \bigoplus_r H^{2r}(W)(-q_r - 1)^{\oplus l_r} & \text{if } n \text{ is even} \end{cases}$$

and $\theta : H^*(W) \rightarrow H^{n+2k-4}(Y)(-1)$ is the surjection in lemma 4.3.

To finish the proof of theorem 1.2 in this case, we need to lift the surjection θ to a surjection onto $H_0^{n+2k-2}(X)$. In order to do that, we construct a section s_U of ψ_2 in geometric way. Following two lemmas will lead us the desired section s_U .

Lemma 4.4. *There is a decomposition*

$$H^{n+2k-4}(Y) \cong \begin{cases} H^{k-2}(\Delta, R^{n+k-2}\pi_*\mathbb{Q}) & \text{if } n \text{ is odd} \\ H^{k-2}(\Delta, R^{n+k-2}\pi_*\mathbb{Q}) \oplus \bigoplus_{p=0, p \neq \frac{k-2}{2}}^{k-2} (\alpha_p \otimes \beta_p)\mathbb{Q} & \text{if } n \text{ is even} \end{cases}$$

where α_p (resp. β_p) is an algebraic cycle generating $H^{2p}(\Delta)$ (resp. $H^{n+2k-4-2p}(\tilde{Q}_t)$).

Proof. Consider the Leray spectral sequence associated to the map $\pi : Y \rightarrow \Delta$.

$$E_2^{pq} = H^p(\Delta, R^q\pi_*\mathbb{Q}) \Rightarrow H^{p+q}(Y, \mathbb{Q})$$

Note that

$$(19) \quad E_2^{pq} = \begin{cases} 0 & \text{if either } q \text{ is odd or } p \text{ is odd, } p = k-2 \\ H^p(\Delta, \mathbb{Q}) & \text{if } p \text{ is even and } q \text{ is even, } q \neq n+k-2 \\ H^p(\Delta, R^{n+k-2}\pi_*\mathbb{Q}) & \text{if } p \text{ is even and } q = n+k-2 \end{cases}$$

where $(R^{n+k}\pi_*\mathbb{Q})_t = \mathbb{Q}^2$ for $t \in \Delta$. Then the spectral sequence degenerates at E_2 . Let $F^\bullet$ be the filtration on $H^{n+2k-4}(Y)$ obtained from the spectral sequence. Then for each $p = 0, 1, \dots, 2(k-2)$ we have a short exact sequence

$$(20) \quad 0 \rightarrow F^{p+1} \rightarrow F^p \rightarrow \text{Gr}_F^p H^{n+2k-4}(Y) \rightarrow 0$$

where

$$\text{Gr}_F^p H^{n+2k-4}(Y) = E_\infty^{p, n+2k-4-p} = H^p(\Delta, R^{n+2k-4-p}\pi_*\mathbb{Q})$$

(Case 1) If n is odd (and hence k is odd): then $n+2k-4$ is odd. Hence $E_2^{pq} = 0$ unless $(p, q) = (k-2, n+k-2)$. Hence we have

$$(21) \quad H^{n+2k-4}(Y) \cong H^{k-2}(\Delta, R^{n+k-2}\pi_*\mathbb{Q})$$

(Case 2) If n is even (and hence k is even): then $n + 2k - 4$ is also even. Note $F^{2p-1} = F^{2p}$ from (19). Hence we may write above sequence as

$$(22) \quad 0 \rightarrow F^{2p+2} \rightarrow F^{2p} \xrightarrow{t_{2p}} \mathrm{Gr}_F^{2p} H^{n+2k-4}(Y) \rightarrow 0$$

First by using descending induction on p , we show that for p such that $k - 2 < 2p \leq 2(k - 2)$ there is a natural splitting of exact sequences (22) such that

$$F^{2p} \cong F^{2p+2} \oplus \mathrm{Gr}_F^{2p} H^{n+2k-4}(Y) \cong \bigoplus_{l=p+1}^{k-2} (\alpha_l \otimes \beta_l) \mathbb{Q}$$

where α_l is an algebraic cycle generating $H^{2l}(\Delta) \cong \mathbb{Q}$ and β_l is an algebraic cycle generating $H^{n+2k-4-2l}(\tilde{Q}_t)(-1) \cong \mathbb{Q}$.

If $2p = 2(k - 2)$, then

$$\begin{aligned} F^{2(k-2)} &\cong \mathrm{Gr}_F^{2(k-2)} H^{n+2k-4}(Y) = E_{\infty}^{2(k-2), n} \\ &= H^{2(k-2)}(\Delta) \otimes H^n(\tilde{Q}_t) \cong (\alpha_{k-2} \otimes \beta_{k-2}) \mathbb{Q} \end{aligned}$$

where α_{k-2} (resp. β_{k-2}) is an algebraic cycle generating $H^{2(k-2)}(\Delta) = \mathbb{Q}$ (resp. $H^n(\tilde{Q}_t) = \mathbb{Q}$). Now suppose $k - 2 < 2p < 2(k - 2)$ and consider the short exact sequence (22). Since $2p > k - 2$, $\dim H^{2p}(\Delta) = \dim H^{n+2k-4-2p}(\tilde{Q}_t) = 1$, hence we can choose an algebraic cycle α_p (resp. β_p) which generates $H^{2p}(\Delta)$ (resp. $H^{n+2k-4-2p}(\tilde{Q}_t)$) and hence

$$\begin{aligned} \mathrm{Gr}_F^{2p} H^{n+2k-4}(Y) &= H^{2p}(\Delta, R^{n+2k-4-2p} \pi_* \mathbb{Q}) = H^{2p}(\Delta) \otimes H^{n+2k-4-2p}(\tilde{Q}_t) \\ &= (\alpha_p \otimes \beta_p) \mathbb{Q} \end{aligned}$$

and by induction hypothesis F^{2p+2} has a decomposition

$$F^{2p+2} \cong \bigoplus_{l=p+1}^{k-2} (\alpha_l \otimes \beta_l) \mathbb{Q}$$

with a basis $\mathcal{B}_{p+1} = \{\alpha_{p+1} \otimes \beta_{p+1}, \dots, \alpha_{k-2} \otimes \beta_{k-2}\}$, where α_l and β_l are algebraic cycles. Let $\mathcal{B}'_p = \{v_p, \alpha_{p+1} \otimes \beta_{p+1}, \dots, \alpha_{k-2} \otimes \beta_{k-2}\}$ be a basis of F^{2p} extending the basis $\mathcal{B}_{p+1}$. Then $t_{2p}(v_p) = q_p(\alpha_p \otimes \beta_p)$ for some nonzero $q_p \in \mathbb{Q}$. Define

$$s_p : \mathrm{Gr}_F^{2p} H^{n+2k-4}(Y) \rightarrow F^{2p}$$

by $s_p(\alpha_p \otimes \beta_p) = \frac{1}{q_p} v_p$. Then s_p is a section of t_p and hence we have a decomposition

$$F^{2p} \cong F^{2p+2} \oplus \mathrm{Gr}_F^{2p} H^{n+2k-4}(Y) = \bigoplus_{l=p+1}^{k-2} (\alpha_l \otimes \beta_l) \mathbb{Q} \oplus s_p(\alpha_p \otimes \beta_p) \mathbb{Q}$$

with a basis $\mathcal{B}_p = \{\alpha_p \otimes \beta_p, \alpha_{p+1} \otimes \beta_{p+1}, \dots, \alpha_{k-2} \otimes \beta_{k-2}\}$ of F^{2p} by identifying $\alpha_p \otimes \beta_p$ with its image under s_p .

Now for splitting for $2p < k - 2$ in the exact sequence (22), the argument is same as the case when $2p > k - 2$, since $\mathrm{Gr}_F^{2p} H^{n+2k-4}(Y) = H^{2p}(\Delta) \otimes H^{n+2k-4-2p}(\tilde{Q}_t) = \mathbb{Q}$. Hence we can choose a natural splitting of each exact sequence (22) and get a decomposition

$$(23) \quad H^{n+2k-4}(Y) \cong H^{k-2}(\Delta, R^{n+k-2} \pi_* \mathbb{Q}) \oplus \bigoplus_{p=0, p \neq \frac{k-2}{2}}^{k-2} (\alpha_p \otimes \beta_p) \mathbb{Q}$$

□

Lemma 4.5. *Let L be the subspace of $H^{n+2k-4}(Y)$ generated by $\{\alpha_p \otimes \beta_p \mid p = 0, \dots, \widehat{\frac{k-2}{2}}, \dots, k-2\}$. Then,*

$$H^{k-2}(\Delta, R^{n+k-2}\pi_*\mathbb{Q}) \cong L^\perp$$

where $\perp$ is the orthogonal complement with respect to the cup product on $H^{n+2k-4}(Y)$.

Proof. First we show that

$$(\alpha_i \otimes \beta_i) \cup (\alpha_j \otimes \beta_j) = \begin{cases} 0 & \text{if } i+j \neq k-2 \\ \deg \Delta & \text{if } i+j = k-2 \end{cases}$$

Since

$$(24) \quad H^{2i}(\Delta, R^{n+2k-4-2i}\pi_*\mathbb{Q}) \otimes H^{2j}(\Delta, R^{n+2k-4-2j}\pi_*\mathbb{Q}) \xrightarrow{\cup} H^{2(i+j)}(\Delta, R^{2n+4k-8-2(i+j)}\pi_*\mathbb{Q})$$

$(\alpha_i \otimes \beta_i) \cup (\alpha_j \otimes \beta_j) = 0$ for $i+j \neq k-2$. For $i+j = k-2$, we observe α_i, β_i closely. Note that for any $i \neq \frac{k-2}{2}$, $\alpha_i = h_\Delta^i$ where $h_\Delta^i \in H^{2i}(\Delta)$ is a class corresponding to an iterated hyperplane section of Δ of codimension i . For β_i , let $\Pi \in H^{n+k-2}(\tilde{Q}_t)$ be a class corresponding to $\mathbb{P}^{\frac{n+k-2}{2}} \subset \tilde{Q}_t$ and $H^r \in H^{2r}(\tilde{Q}_t)$ a class corresponding to an iterated hyperplane section of $\tilde{Q}_t$ of codimension r . Note that Π can be chosen in either families F_t^1 or F_t^2 of $\mathbb{P}^{\frac{n+k-2}{2}}$ contained in $\tilde{Q}_t$ ([R]). Then,

$$\beta_i = \begin{cases} H^{\frac{n+2k-4-2i}{2}} & \text{if } i > \frac{k-2}{2} \\ \Pi \cup H^{\frac{k-2}{2}-i} & \text{if } i < \frac{k-2}{2} \end{cases}$$

Then for $i+j = k+2$,

$$(\alpha_i \otimes \beta_i) \cup (\alpha_j \otimes \beta_j) = \deg \Delta$$

Now we show $L \cap L^\perp = 0$. Let $\eta = \sum_{i=0, i \neq \frac{k-2}{2}}^{k-2} c_i (\alpha_i \otimes \beta_i) \in L \cap L^\perp$. Then, we have $c_i \cdot \deg \Delta = 0$ for any $i \neq \frac{k-2}{2}$. Hence $\eta = 0$. Hence

$$H^{n+2k-4}(Y) = L \oplus H^{k-2}(\Delta, R^{n+k-2}\pi_*\mathbb{Q}) = L \oplus L^\perp$$

Hence we get

$$H^{k-2}(\Delta, R^{n+k-2}\pi_*\mathbb{Q}) = L^\perp$$

□

From the above lemmas, we have an injection

$$(25) \quad h : H^{k-2}(\Delta, R^{n+k}p_{1*}\mathbb{Q}) \hookrightarrow H^{n+2k-2}(X_\Delta)$$

We refer the diagram (18) for the following lemma.

Lemma 4.6. *There is a section defined in geometric way*

$$s_U : Gr_{n+2k-2}^W H_c^{n+2k-1}(X_U) \rightarrow H^{n+2k-4}(Y)(-1)$$

of ψ_2 .

Proof. First recall $H^{n+2k-4}(Y)(-1) \cong H^{n+2k-2}(X_\Delta)$ and ψ_2 is the connecting homomorphism in the exact sequence of mixed Hodge structures (14)

$$\dots \rightarrow H^{n+2k-2}(X) \rightarrow H^{n+2k-2}(X_\Delta) \xrightarrow{\psi_2} Gr_{n+2k-2}^W H_c^{n+2k-1}(X_U) \rightarrow 0$$

As in the proof of lemma 4.2, we can show that the the Leray spectral sequence associated to $p_1 : X_U \rightarrow U$ gives rise to an exact sequence

$$0 \rightarrow E_\infty^{2k-2, n+1} \rightarrow H_c^{n+2k-1}(X_U) \rightarrow E_\infty^{k-1, n+k} \rightarrow 0$$

By taking the exact functor Gr_{n+2k-2}^W , we get an isomorphism

$$l : \mathrm{Gr}_{n+2k-2}^W H^{n+2k-1}(X_U) \xrightarrow{\cong} \mathrm{Gr}_{k-2}^W H_c^{k-1}(U)$$

since $E_\infty^{2k-2, n+1} = H_c^{2k-2}(U)(-\frac{n+1}{2}) \cong H^{2k-2}(\mathbb{P}^{k-1})(-\frac{n+1}{2})$ is a pure Hodge structure of weight $n+2k-2$, if n is odd and 0 if n is even. Hence we have a following commutative diagram:

$$(26) \quad \begin{array}{ccc} H^{n+2k-2}(X_\Delta) & \xrightarrow{\psi_2} & \mathrm{Gr}_{n+2k-2}^W H_c^{n+2k-1}(X_U) \\ \uparrow h & & \downarrow l \cong \\ H^{k-2}(\Delta, R^{n+k} p_{1*} \mathbb{Q}) & \xrightarrow{\delta} & \mathrm{Gr}_{k-2}^W H_c^{k-1}(U)(-\frac{n+k}{2}) \end{array}$$

where h is the injection of (25) and $\delta : H^{k-2}(\Delta, R^{n+k} p_{1*} \mathbb{Q}) \rightarrow \mathrm{Gr}_{k-2}^W H_c^{k-1}(U)(-\frac{n+k}{2})$ is the connecting homomorphism of long exact sequence induced by a short exact sequence of sheaves on $\mathbb{P}^{k-1}$

$$0 \rightarrow j_!(R^{n+k} p_1 \mathbb{Q})|_U \rightarrow R^{n+k} p_{1*} \mathbb{Q} \rightarrow i_{\Delta*}(R^{n+k} p_{1*} \mathbb{Q})|_\Delta \rightarrow 0$$

where $j : U \rightarrow \mathbb{P}^{k-1}$ and $i_\Delta : \Delta \rightarrow \mathbb{P}^{k-1}$ are inclusions. Hence in order to choose a section of ψ_2 , it is enough to construct a section of δ .

Note that $(R^{n+k} p_{1*} \mathbb{Q})_t = \mathbb{Q}^2$ for $t \in \Delta$ and $R^{n+k} p_{1*} \mathbb{Q}|_U = \mathbb{Q}_U$ is the constant sheaf. Let G be the monodromy group of $R^{n+k} p_{1*} \mathbb{Q}|_\Delta$. Then $R^{n+k} p_{1*} \mathbb{Q}|_\Delta^G = \mathbb{Q}$ is a subsheaf of $R^{n+k} p_{1*} \mathbb{Q}|_\Delta$ and let $R^{n+k} p_{1*} \mathbb{Q}|_\Delta^G \hookrightarrow R^{n+k} p_{1*} \mathbb{Q}|_\Delta$ be an injection. Let

$$s : i_{\Delta*}(R^{n+k} p_{1*} \mathbb{Q}|_\Delta^G) \rightarrow i_{\Delta*}(R^{n+k} p_{1*} \mathbb{Q}|_\Delta)$$

be the induced map. Then there is a sheaf $s^*(R^{n+k} p_{1*} \mathbb{Q})$ on $\mathbb{P}^{k-1}$ which fits into the following commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & j_!(R^{n+k} p_1 \mathbb{Q}|_U) & \longrightarrow & s^*(R^{n+k} p_{1*} \mathbb{Q}) & \longrightarrow & i_{\Delta*}(R^{n+k} p_{1*} \mathbb{Q}|_\Delta^G) \rightarrow 0 \\ & & \parallel & & \downarrow & & \downarrow s \\ 0 & \longrightarrow & j_!(R^{n+k} p_1 \mathbb{Q}|_U) & \longrightarrow & R^{n+k} p_{1*} \mathbb{Q} & \longrightarrow & i_{\Delta*}(R^{n+k} p_{1*} \mathbb{Q}|_\Delta) \rightarrow 0 \end{array}$$

From this, we get a commutative diagram

$$\begin{array}{ccccccc} \dots H^{k-2}(\mathbb{P}^{k-1}, s^*(R^{n+k} p_{1*} \mathbb{Q})) & \longrightarrow & H^{k-2}(\Delta, \mathbb{Q}) & \xrightarrow{\delta_1} & \mathrm{Gr}_{k-2}^W H_c^{k-1}(U, \mathbb{Q}) & \longrightarrow & 0 \\ \downarrow & & \downarrow s & & \parallel & & \\ \dots H^{k-2}(\mathbb{P}^{k-1}, R^{n+k} p_{1*} \mathbb{Q}) & \longrightarrow & H^{k-2}(\Delta, R^{n+k} p_{1*} \mathbb{Q}) & \xrightarrow{\delta} & \mathrm{Gr}_{k-2}^W H_c^{k-1}(U) & \longrightarrow & 0 \end{array}$$

i.e.

$$(27) \quad \delta_1 = \delta \circ s$$

Now from the exact sequence

$$\dots \rightarrow H^{k-2}(\mathbb{P}^{k-2}) \xrightarrow{i_\Delta^*} H^{k-2}(\Delta) \xrightarrow{\delta_1} \mathrm{Gr}_{k-2}^W H_c^{k-1}(U) \rightarrow 0$$

we have an isomorphism

$$Gr_{k-2}^W H_c^{k-1}(U) \xrightarrow{\cong} H^{k-2}(\Delta)/\text{im } i_\Delta^*$$

hence we can choose a natural section $s_1 : Gr_{k-2}^W H_c^{k-1}(U) \rightarrow H^{k-2}(\Delta)$ of $\delta_1 : H^{k-2}(\Delta) \rightarrow Gr_{k-2}^W H_c^{k-1}(U)$, i.e. $\delta_1 \circ s_1 = \text{id}$. Then from (27) we get a section $s \circ s_1$ of δ . By combining all these, we get a commutative diagram:

$$\begin{array}{ccc} H^{n+2k-2}(X_\Delta) & \xrightarrow{\psi_2} & Gr_{n+2k-2}^W H_c^{n+2k-1}(X_U) \\ \uparrow h & & \downarrow l \cong \\ H^{k-2}(\Delta, R^{n+k} p_{1*} \mathbb{Q}) & \xrightarrow{\delta} & Gr_{k-2}^W H_c^{k-1}(U) \left(-\frac{n+k}{2}\right) \\ \uparrow s & & \parallel \\ H^{k-2}(\Delta) & \xrightarrow{\delta_1} & Gr_{k-2}^W H_c^{k-1}(U) \left(-\frac{n+k}{2}\right) \\ & \xleftarrow{s_1} & \end{array}$$

Set

$$s_U = h \circ s \circ s_1 \circ l$$

then

$$\psi_2 \circ s_U = \psi_2 \circ (h \circ s \circ s_1 \circ l) = l^{-1} \circ (\delta \circ s \circ s_1) \circ l = l^{-1} \circ l = \text{id}$$

Hence s_U is a section of ψ_2 . □

Proof of theorem 1.2 when $n+k$ is even. Now we finish the proof of the Theorem 1.2 in case when $n+k$ is even. Recall the diagram (18)

$$\begin{array}{ccccccc} & & H^*(W) & & & & \\ & & \downarrow \theta & \searrow \psi_2 \circ \theta & & & \\ 0 \rightarrow & H_0^{n+2k-2}(X) & \xrightarrow{\psi_1} & H^{n+2k-4}(Y)(-1) & \xrightarrow{\psi_2} & Gr_{n+2k-2}^W H_c^{n+2k-1}(X_U) & \rightarrow \dots \\ & & & & \xleftarrow{s_U} & & \end{array}$$

where s_U is the section of ψ_2 constructed in lemma 4.6.

Let

$$\Theta = \theta - s_U \circ \psi_2 \circ \theta : H^*(W) \rightarrow H^{n+2k-4}(Y)(-1)$$

First note that $\psi_2 \circ \Theta = 0$: In fact,

$$\begin{aligned} \psi_2 \circ \Theta &= \psi_2 \circ (\theta - s_U \circ \psi_2 \circ \theta) \\ &= \psi_2 \circ \theta - \psi_2 \circ s_U \circ \psi_2 \circ \theta = \psi_2 \circ \theta - \psi_2 \circ \theta = 0 \end{aligned}$$

Hence $\text{im } \Theta \subseteq \ker \psi_2$ and we may consider that Θ is mapped into $H_0^{n+2k-2}(X)$ since ψ_1 is an injection.

$$\begin{array}{ccccccc} & & H^*(W) & & & & \\ & \swarrow \Theta = \theta - s_U \circ \psi_2 \circ \theta & \downarrow \theta & \searrow \psi_2 \circ \theta & & & \\ 0 \rightarrow & H_0^{n+2k-2}(X) & \xrightarrow{\psi_1} & H^{n+2k-4}(Y)(-1) & \xrightarrow{\psi_2} & Gr_{n+2k-2}^W H_c^{n+2k-1}(X_U) & \rightarrow \dots \\ & & & & \xleftarrow{s_U} & & \end{array}$$

Now we show this map is in fact a surjection. Let $\alpha \in H_0^{n+2k-2}(X)$. Since ψ_1 is an injection we may identify α with $\psi_1(\alpha)$. Then there is $\beta \in H^*(W)$ such that $\theta(\beta) = \psi_1(\alpha)$. Then

$$\begin{aligned}\Theta(\beta) &= (\theta - s_U \circ \psi_2 \circ \theta)(\beta) \\ &= \theta(\beta) - s_U \circ \psi_2 \circ \theta(\beta) = \psi_1(\alpha) - s_U \circ \psi_2 \circ \psi_1(\alpha) = \psi_1(\alpha) = \alpha\end{aligned}$$

Hence we have a surjection

$$\Theta : H^*(W) \longrightarrow H_0^{n+2k-2}(X)$$

as we claimed. This finishes the proof of theorem 1.2. $\square$

REFERENCES

- [A] D. Arapura, *The Leray spectral sequence is motivic*, Invent. math. 160. (2005) 567 – 589
- [E] C. Ehresmann, *Sur la topologie de Certains Espaces Homogenes*, Annals of Math. 2nd Ser., Vol 35, No 2. (1934) 396 – 443
- [F] W. Fulton, *Intersection Theory*, 2nd ed., Springer-Verlag, Berlin, 1998.
- [G] A. Grothendieck, *Hodge's general conjecture is false for trivial reasons*, Topology 8, (1969)
- [GH] P. Griffiths, J. Harris, *Principles of Algebraic geometry*, John Wiley & Sons, 1978
- [L] J. Lewis, *A survey of the Hodge conjecture* CRM Monograph Series, 10. American Mathematical Society, (1999).
- [O] K. O'Grady, *The Hodge structure of the Intersection of Three quadrics in an odd dimensional projective space*, Math. Ann. 273, (1986) 277 – 285
- [R] M. Reid, *The complete intersection of two or more quadrics*, Ph.D. thesis (1972)
- [S] Spanier, *algebraic Topology*, McGraw-Hill, 1966

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, WEST LAFAYETTE, IN 47907, U.S.A.
E-mail address: `sjkang@math.purdue.edu`